

# Isolation and Enzymatic Activity of Thermophilic Bacteria from Waste Dump Sites in Umudike Town

Obi, C. N\*, Ihuoma, B. O. and Ike, S.

Department of Microbiology, College of Natural Sciences, Michael Okpara University of Agriculture, Umudike, P.M.B 7267 Umuahia, Abia State Nigeria.

## ARTICLE INFO

### Article history:

Received: 24 May 2021;

Received in revised form:

19 June 2021;

Accepted: 30 June 2021;

### Keywords

Activity,  
Bacteria,  
Enzyme,  
pH,  
Thermophilic Temperature.

## ABSTRACT

Enzymatic potentials of thermo-tolerant bacteria from waste dumpsites were analyzed. 30 soil samples were collected using sterile spatular into sterile universal bottles, labelled and taken to the laboratory analyses. The soil samples were serially diluted and inoculated by spread plate method on different media and incubated at 30°C for 48 hrs. The pure isolates were stored in agar slants in the refrigerator for further use. Isolates with thermophilic ability were screened for amylase, protease and lipase activity. Effects of temperature and pH on enzyme activity of the thermophilic bacterial isolates were also assessed. *Bacillus subtilis* (70.0%), *Bacillus licheniformis* (43.3%), *Bacillus cereus* (56.7%), *Pseudomonas aeruginosa* (56.7%), *Streptomyces* species (36.7%), *Bacillus brevis* (43.3%) and *Nocardia* species (30%) were isolated from the soil samples. All the bacterial isolates tolerated 55°C temperature while only *Bacillus cereus*, *Bacillus brevis* and *Nocardia* species were isolated at 60°C. All the isolates except *Pseudomonas aeruginosa* gave amylase activity while all the isolates gave protease activity. All the isolates except *Bacillus licheniformis* and *Nocardia species* showed lipase activity. Temperature and pH had varied effects on the enzyme activity of the isolates. These bacteria are good sources of enzymes at thermophilic temperatures and varied pH for commercial use.

© 2021 Elixir All rights reserved.

## 1.0 Introduction

Thermophilic bacteria are microbes that mostly inhabit hot environments, live and survive in temperatures above 70°C (Decastro *et al.*, 2016). They have been less explored due to difficulties in isolation and maintenance of pure culture. Therefore, their diversity and biotechnological potential remains to be explored from majority of the thermal habitats. The adaptation to these harsh habitats explains the high genomic and metabolic flexibility of microbial communities in these ecosystems and makes thermophiles and their thermostable proteins very suitable for some industrial and biotechnological applications (Decastro *et al.*, 2016). As a consequence of growth at high temperatures and unique macromolecular properties, thermophiles can possess high metabolism, physically and chemically stable enzymes and lower growth but higher end product yields than similar mesophilic species (Lewin *et al.*, 2013).

Waste dump can be defined as waste type that includes predominantly household waste (domestic waste) with sometimes the addition of commercial waste collected by a municipality and deposited within a given area (Odeyemi, 2012).

Thermostable proteases are of greater advantage in applications because they not only do not usually denature at high temperatures, but they also remain active at such temperatures. Several workers have reported thermophilic bacteria from diverse environmental habitats such as geothermal sites and hot springs. Protease constitute a large and complex group of hydrolytic enzymes with important

application in medical, pharmaceutical, biotechnology, leather, detergents and food industries (Ramakrishna *et al.*, 2010).

Microorganisms present a remarkable potential for proteolytic enzymes production due to their extensive biochemical diversity and susceptibility to genetic manipulation (Soares and Sato, 2014).

Amylases are enzymes that break down starch or glycogen. Recent discoveries of starch degrading enzymes have led to increased application of amylases in various industrial processes. Thermostable amylase not only solves stability issues, but also accelerates reaction rate, reduces possible contaminations and lowers viscosity of medium, thus directly benefitting the starch-processing industries under high temperatures (Homaei *et al.*, 2016).

Lipase are serine hydrolase that catalyze both hydrolysis and occupy a place of prominence among biocatalysts owing to their ability to catalyze a wide range of reactions and are an important group of biotechnologically relevant enzymes and they find a massive applications (Mohammed 2013).

Many industries need thermophiles for the production of thermostable enzymes that have wide applications, hence there is a need for isolation of specific bacteria that will be of economic importance.

This research aimed at isolation and identification of thermophilic bacteria from soil waste dump site samples with ability to produce industrial enzymes.

## 2.0 Materials and Methods

### 2.1 Sample collection

Thirty (30) soil samples were collected from 6 different sites within Umudike Metropolis; Ahiaeke, Government College, Timber, NDDC GEJ Hostels of Michael Okpara University of Agriculture as well as Living Rock Church Junction respectively in sterile universal bottles using a sterile soil spatula and immediately taken to the laboratory for analysis.

### 2.2 Cultivation and enumeration of total bacterial population

One gram of each soil sample was diluted in 9ml of sterile peptone water in 250ml conical flask and placed in an orbital shaker at 150 rpm for 15 minutes to get a homogenized soil suspension. 0.1 ml of the diluted soil sample was aseptically inoculated onto sterile Petri dishes containing Nutrient Agar (HIMEDIA, India) containing cycloheximide to inhibit contaminating fungi in triplicates and incubated in Southern Biological Incubator at 37°C for 24 hours. After incubation, the number of colonies on each plate was counted using a colony counter and the average count determined (Mrunmaya *et al.*, 2012). The isolates were sub-cultured to get pure isolates which were later Gram Stained, subjected to biochemical and sugar fermentations tests.

### 2.3 Determination of bacterial thermo-tolerance

Pure cultures of 24 hour old of the bacterial isolates were tested for their thermo-tolerant ability by inoculating them into 5 ml of Nutrient broth medium (HIMEDIA) in test tubes. The tubes were incubated at 45°C for 12 hours after which each broth culture was streaked onto freshly prepared Nutrient Agar medium and incubated at 37°C for 24 hr to determine their viability (Mrunmaya *et al.*, 2012).

### 2.4 Determination of enzyme activity

#### 2.4.1 Amylase activity

Starch hydrolysis method was used to identify the amylolytic properties of the thermotolerant isolates. The isolates were streaked on the Starch Agar Plates and incubated at 37°C for 24 hours. After incubation, 1% iodine solution was flooded on the starch agar plate. The presence of blue color around the growth indicated negative results (Cowan, 2016).

#### 2.4.2 Protease activity

Protease activity was detected on Muller-Hinton Agar (HIMEDIA) containing 3% Bovine Serum Albumen (BSA). The plates were streaked with test isolates and incubated at 37°C for 24 hr. The presence of a transparent zone around the colonies indicated caseinase activity (Burke *et al.*, 2015).

Lipase activity was determined on tributyrine agar streaked with bacterial isolates and incubated immediately at 37°C for 24-48 hours. Plates was observed for clear zone around the bacterial growth (Cappuccinno, 2008)

### 2.5 Effect of temperature on enzyme activities

The isolates were streaked on the starch agar plates and incubated at 30°C, 35°C, 40°C, 45°C, 50°C, 55°C and 60°C respectively for 24 hours. After incubation, 1% iodine solution was flooded on the starch agar plate. The presence of brown color around the growth indicated negative results while blue-black colour showed positive result (Cowan, 2016).

### 2.6 Effect of pH on enzyme Activities

The medium for each enzymes was prepared and six buffer solutions with pH of 3.0, 4.0, 5.0, 6.0, 7.0, 8.0 were labeled in six beakers in order to differentiate the six different buffers. The pH for each enzyme were initially tested using the pH meter and adjusted by adding 5ml of appropriate buffer into the respective beakers using the graduated pipette. After the appropriate pH was determined, the test organism was added and incubated at 37°C for 24 hours (Campbell *et al.*, 2011).

## 3.0 Results

Table 1 shows the bacterial isolates from the waste dumpsites which are *Bacillus subtilis*, *B. cereus*, *B. brevis*, *B. licheniformis*, *Pseudomonas aeruginosa*, *Nocardia* species and *Streptomyces* species.

Fig. 1 shows the occurrence of bacteria in the soil samples. *Bacillus subtilis* had the highest percentage occurrence (70) while *Nocardia* species had the least (30%).




Fig 1. Frequency of occurrence of bacterial isolates from dump sites.

Table 2 shows the thermo-tolerance result. All the bacteria had profuse growth at 30°C, but moderate growth at 45°C. At 50°C and 55°C respectively, all the bacteria had scanty growth, but 60°C, only *B. cereus*, *B. brevis* and *Nocardia* species grew though scantily.

Table 1. Morphological and biochemical characteristics of the bacterial isolates.

Colony	Grams reaction	Spore	Flagella	Motility	Cell shape	Catalase	Indole	Oxidase	Mannitol	Sucrose	Fructose	Lactose	Suspected organism
Circular, rough, white	+	+	+	+	Short chains	+	-	+	A/-	A/-	-	-	<i>Bacillus subtilis</i>
White, raised	+	+	+	+	Irregular	+	-	+	A/-	A/-	+	A/-	<i>Bacillus cereus</i>
Pale yellow	+	+	+	+	Rods	+	-	+	A/-	A/-	-	-	<i>Bacillus brevis</i>
Rough	+	+	+	+	Convex irregular	+	-	-	+	+	+	+	<i>Bacillus licheniformis</i>
Capsulated	-	Non spore	single	+	rod	+	-	+	+	-	-	-	<i>Pseudomonas aeruginosa</i>
Non capsulated	+	-	-	-	rod	+	-	+	-	-	-	-	<i>Nocardia</i> species
filamentous	+	+	+	-	chain	+	-	+	-	-	+	-	<i>Streptomyces</i> species

KEY: + = Positive - = negative, A = acid present

**Table 2. Thermo-tolerance of bacterial isolates.**

Bacterial isolates	30°C	35°C	40°C	45°C	50°C	55°C	60°C
<i>Bacillus subtilis</i>	+++	+++	+++	++	+	+	-
<i>Bacillus cereus</i>	+++	+++	+++	++	+	+	+
<i>Bacillus licheniformis</i>	+++	+++	+++	++	+	+	-
<i>Pseudomonas aeruginosa</i>	+++	+++	+++	++	+	+	-
<i>Streptomyces</i> species	+++	+++	+++	++	+	+	-
<i>Nocardia</i> species	+++	+++	+++	++	+	+	+
<i>Bacillus brevis</i>	+++	+++	+++	++	+	+	++

**Key:**

+++ = Profuse positive growth

++ = Moderate positive growth

+ = Scanty growth

- = No growth

Table 3 shows that all the bacterial isolates except *Pseudomonas aeruginosa* had amylase activity while all the isolates had protease activity. All the isolates had lipase activity except *B. licheniformis* and *Nocardia* species.

**Table 3. Enzymatic activity of the bacterial isolates.**

Bacterial isolates	Amylase	Protease	Lipase
<i>Bacillus subtilis</i>	+	+	+
<i>Bacillus brevis</i>	+	+	+
<i>Bacillus cereus</i>	+	+	+
<i>Bacillus licheniformis</i>	+	+	-
<i>Pseudomonas aeruginosa</i>	-	+	+
<i>Nocardia</i> species	+	+	-
<i>Streptomyces</i> species	+	+	+

**Key:**

+ = positive

- = negative

Fig. 2 shows the effect of temperature on bacterial enzyme activity. The activity was optimum at 30°C, but gradually decreased with increase in temperature up to 60°C. The highest activity at 60°C was from *B. licheniformis* while the least activity was from *Streptomyces* species.

Effect of pH on the bacterial enzyme activity is shown on Fig. 3. Generally, there was an increase in activity from pH 3.0 to 7.0 with best activity recorded at pH 7 for all the isolates except *Nocardia* species whose best activity was at pH 6. There was a general decrease in activity beyond pH 7.0

**4.0 Discussion**

The present research aimed at isolation of bacteria from waste dumpsites with enzymatic activities. Based on morphological and biochemical characterization, the dumpsites were found to harbour a variety of temperature tolerant and enzyme producing bacterial species which include *Bacillus subtilis*, *B. cereus*, *B. brevis*, *B. licheniformis*, *Pseudomonas aeruginosa*, *Nocardia* species and *Streptomyces* species. The isolates recovered from the waste dumpsites are in agreement with the findings of Obianefo *et al.* (2017) who worked on dumpsites in Port Harcourt, South South Nigeria.

The occurrence of bacteria isolates have been reported to be associated with biodegradation of raw human faecal discharges and other wastes at the dumpsites. This observation compares favorably with the report of Hammond and Belies (1986). The micro flora of habitats is determined by the factors which include the availability of nutrients and suitable physical conditions of moisture, temperature etc. Microorganisms that produce amylases could be isolated from places such as soils around rice mills, cassava farms and flour markets (Fossi *et al.*, 2005). Vasekaran *et al.* (2010) stated

that amylase producing bacterial strains were isolated from the soils contaminated with decaying materials including kitchen waste, bakery waste, flour mill waste, soil receiving tea waste, hot white rice gruel and compost heap.

The ability of bacteria to grow at high temperatures is indicative of their potentials to be used at industrial level application. The application of microbial based enzymes at high temperature is desirable in the food industries where thermal treatments are applicable for the control of industrial microorganisms in the product (Cowan, 1996). Considering these facts, the thermophilic bacteria were found to possess amylase, protease and lipase activity. Rath *et al.* (2010) have also reported various extracellular enzymes from some thermophilic bacteria isolated from the same hot spring. However, several workers have reported protease activity from thermophilic *Bacillus* species (Guangrong *et al.*, 2006 and Vijayalakshmi *et al.*, 2011).

Proteases are essential constituents of all forms of life on earth including prokaryotes, fungi, plants and animals and are highly exploited enzymes in food, leather, detergent, pharmaceutical, diagnostics, waste management and silver recovery (Babu and Lakshmi, 2005). *Bacillus*' protease is of special importance because of its wide applications in various industries like pharmaceutical, leather, food and waste processing industries (Pastor *et al.*, 2001).

The variations in the optimum temperature for the production of different enzymes as recorded in this work is attributed to genetics of the different isolates. Reports have shown that temperature affects both the production and activities of enzymes. Notwithstanding, organisms can be very active over a wide temperature ranges as thermophiles and extreme thermophiles produce enzymes of industrial purposes. The findings here show that these organisms produce enzymes at elevated temperatures (50°C, 55°C and 60°C; Table 2) and as such can be useful for commercial use. Also the ability to produce enzyme at such temperature could be utilized in the control of contaminations by microorganisms.

Lipase production by the organisms imply potential use in both food industry and laundry for the removal of strong oil stains on fabrics. Furthermore, the results show that there were variations in the optimum pH for the production of the different enzyme types (amylase, protease and lipase). *Bacillus cereus* and *subtilis* had optimum pH for amylase production at pH 6.0 while the optimum pH for *Bacillus brevis* and *Bacillus licheniformis* was 7.0 and both had exactly the same rate of amylase production (0.0318µmol/min) at this pH.




Fig 2. Effect of Temperature on Bacterial Enzyme Activity




Fig 3. Effect of pH on Bacterial Enzyme Activities.

*Pseudomonas aeruginosa* produced amylase optimally at pH 7.0 with production rate of 0.0258 $\mu$ mol/min. The optimum pH for amylase by *Nocardia* species and *Streptomyces* species isolates was 6.0 for both while their rate of enzyme production were 0.0280 $\mu$ mol/min and 0.0298 $\mu$ mol/min respectively.

Protease production rate varied among the bacteria isolate in terms of optimum pH and rate of production. The optimum pH for protease production was 6.0 for four bacteria isolates: *Bacillus cereus*, *Bacillus licheniformis*, *Nocardia* species and *Streptomyces* species. However their rates of production varied (Fig. 3). *Bacillus brevis* and *Pseudomonas aeruginosa* recorded optimum protease at pH of 7.0 with the production rates of 0.0212 $\mu$ mol/min and 0.0204 $\mu$ mol/min respectively. This agrees with the findings of Mrunmaya (2013).

The rate of Lipase production by the bacterial isolates was comparatively lower than those of amylase and protease productions. The variations in the enzyme production cut across the different organisms for a particular type of enzyme as well as variation between the same organisms for different enzyme types.

Some enzymes were active over a range of pH values. Vihinen *et al.* (1989) reported that some enzymes are active and stable over a wide range of pH 4.0 to 11.0. However, extreme pH values have been reported to inhibit enzyme activity.

### Conclusion

The results obtained in this project work showed that dump site soil harboured a wide range of bacteria which were confirmed to be thermophiles. Most of the bacterial isolates demonstrated ability to produce enzymes including amylase, protease and lipase at elevated temperatures. Results also showed variations in the optimum conditions for enzyme production. The optimum pH and temperature of enzymes production reveals that the isolates have potentials as source of different enzymes that could be of various industrial benefits.

### Recommendations

Waste dumpsites are good sources of industrially beneficial bacterial that could be assessed for various industrial applications. The search for such organisms from waste dumpsites could furnish scientists and industrialists with better performing bacteria in the future.

### References

DeCastro, M. E., Rodriguez-Belmonte E., Gonzalez-siso M.I (2016) .Metagenomics of Thermophiles with a Focus on Discovery of Novel Thermozymses. *Front. Microbial.*,7: 15-21.

Lewin, A., Wentzel, A. and Valla, S. (2013). Metagenomics of microbial life in extreme temperature environments. *Current Opinion in Biotechnology* 24(3).

Odeyemi, A.T. (2012). Antibioqram status of bacterial isolates from air around dump site of Ekiti State Destitute Centre at Ilokun, Ado-Ekiti, Nigeria. *Journal of Microbiology Research*, 2(2):12-18.

Ramakrishna, D. P. N., Gopi Reddy, N. and Rajagopal, S. V. (2010). Purification and properties of an extra cellular alkaline protease produced by *Bacillus subtilis*. *Int. J. Biotechnol. Biochem.* 6(4): 493-504.

Soares de Castro, R. J. and Sato, H. H. (2014). "Production and biochemical characterization of protease from *Aspergillus oryzae*: an evaluation of the physical-chemical parameters using agroindustrial wastes as supports," *Biocatalysis and Agricultural Biotechnology*, (3)3: 20–25.

Homaei, A., Ghanbarzadeh, M. and Monsef, F. (2016). "Biochemical features and kinetic properties of alpha amylases from marine organisms," *International Journal of Biological Macromolecules*, 83: 306-314.

Mohammed, H. J. (2013) Lipase activity of *Acinetobacter baumannii* 82 Physicochemical Factors Affecting the local hot spring Tarabalo, Nayagarh District, Orissa, India. *Internet J Microbiol.* 7: 2.

Mrunmaya, K. P., Chandi, C. R. and Kumananda, T. (2012). Bioactivities of bacterial endophytes isolated from leaf tissues of *Hyptis suaveolens* against some clinically significant pathogens. *Journal of Applied Pharmaceutical Science*, 7 (08), 131-136.

Cowan, D. (1996). Industrial enzyme technology. *Trends Biotechnol.* 14: 177-178.

Burke, V., Robinson, J. O., Richardson, C. J. L., and Bundell, C. S. (2015). Longitudinal studies of virulence factors of *pseudomonas aeruginosa* in cystic fibrosis. *Pathology*, (23)2: 145– 148.

Cappuccino J. G. and Sherman N. (2008). Microbiology: A Laboratory Manual, 8<sup>th</sup> edition. Pearson benjamin Cummings, San Francisco, CA, USA.

Campbell, N.A., Reece, J.B., Urry, L.A., Cain, M.L., Wasserman, S.A., Minorsky, P.V., Jackson, R.B. (2011).

Obianefo, F. U; Chindah, A. C; Ochekwu, E. B (2017). Water quality and phytoplankton distribution Nta- Wogba stream receiving municipal discharges in Port Harcourt, Rivers state Nigeria. *Research journal of Environmental Toxicology*, 10:135-143.

Hammond, P.B. and Belles, R. P. (1986). Waste Management in the Coastal Areas of the ASEAN Region

Karnwal, A and V. Nigam, (2013) Production of amylase enzyme by isolated microorganisms and its application. *IJPBS*, 3(4): 354-360.

Vaseekaran, S., Balakumar, S and Arasaratnam, V (2010) Isolation and identification of a bacterial strain producing thermostable Amylase, *Tropical Agricultu. Resear.*, 22(1): 1-11

Rath, C.C., (2010). Heat stable lipase activity of thermotolerant bacteria from hot springs at Orissa, India. *Cytobios.* 99: 105-11.

Guangrong H., Tiejing Y., Po H., Jiaying J. (2006). Purification and characterization of a protease from Thermophilic *Bacillus* strain HS08. *Afr J Biotechnol.* 5: 2433-2438.

Vijayalakshmi, S., Venkatkumar, S., Thankamani V., (2011). Screening of alkalophilic thermophilic protease isolated from *Bacillus* RV.B2.90 for industrial applications. *Res Biotechnol.* 2: 32.

Babu, N.K.S., Lakshmi, K.D. (2005). Optimization of thermostable alkaline protease production from species of *Bacillus* using rice bran. *Afr J Biotechnol.* 4: 724-726

Pastor, MD., Lorda, G.S and Balatti. A. (2001). Protease obtention using *Bacillus subtilis* 3411 and amaranth seed meal medium at different aeration rates. *Braz J Microbiol.* 32: 1-8.

Padhiar, A.R and S. Kommu, 2016. Isolation, characterization and optimization of bacteria producing amylase. *Int. J. Adv. Res. Biol. Sci.*, 3(7): 1-7. Vihinen, M and Mantsala, P (1989). Microbial amyolytic enzyme. *Crit.. biochemical. Molecular. Biology.* 24: 329 – 418.

**APPENDIX**



**Positive Amylase Test**