



Study of cladocera diversity with reference to chydoridae and bosmanidae family of nira left bank canal baramati and Tarangawadi Lake of indapur Taluka district Pune

U.S.Ghantaloo^{1,*}, J.P.Sarwade² and S.M.Kamble³

¹Shardabai Pawar Mahila College Baramati, Dr Babasaheb Ambedkar Marthwada University

²Arts Commerce & Science College, Indapur.

³Zoology Department, S.D.M.M. College, Kalamb.

ARTICLE INFO

Article history:

Received: 14 June 2012;

Received in revised form:

15 November 2015;

Accepted: 20 November 2015;

Keywords

Labrum,
Antennules,
Dorsal,
Ventral,
Post abdominal claw.

ABSTRACT

The present paper focuses on the study of cladocera of nira left bank canal and Tarangawadi lake of indapur and baramati region cladocera are fresh water zooplankton inhabiting all the niches of fresh water bodies. The study of cladocera has being fascinating subject to the biologist. In the digestive tract of most fishes one to ninety percent cladocera are found. Hence in order to increase pisciculture study of cladocerans are important. The water samples were collected between 8 am to 11am during the year Dec 2010 to august 2011. The collected samples were preserved in 4% formalin to study diversity of chydoridae family. The species identified by standard key method. About 17 species have been recorded identification was based on the presence of body parts

© 2015 Elixir All rights reserved.

Introduction

Cladocerans are fresh water zooplankton inhabiting all the niches of fresh water bodies. The cladocerans has been the object of microscopic study dating back to 18th century. Cladocerans are minute forms the average length being 0.2mm to 3.5mm.

Swammerdan (1969), Muller (1785), Daday(1889) and sars (1901) revealed the valuable information about the cladocerans. The cladocerans fauna of the united states has been the result of several workers. Birge (1918), Pennak (1953) and Brook (1959) wrote a comprehensive chapter on the cladocerans of the united state. The book on chydoridae cladocerans of the ussr is a notable contribution from smirnov (1974) Near about 110 species have been recorded from india (patil and Goundar 1984).

The cladocerans are divided into 7 families Daphnidae, Chydoridae, Macrothriadae, Moinidae are found in india and Polyphenidae, Leptidoridae, Holopodidae are characteristically absent in the inland water of India. Family Chydoridae is divided into orders Alona, Biapertura, pleuroxus, dunhevida, chydorus.

Topography of Area


The water bodies of Baramati and indapur region were selected for water sampling. Nira left bank canal is built on the veer dam and flows through the purandar, baramati and indapur region it lies at latitude 18° 08' 28" N and longitude 74° 32' 02" E Tarangawadi lake is situated in indapur region it is 50 km away from baramati region water from Khadakwasala dam water is stored in the lake it lies at latitude about 18° N and 19° N Latitude and 75° E longitude East.

Material Method

The water sample were collected from both Nira left bank canal baramati region and Tarangawadi lake of indapur region. The samples were collected using plankton net and stored in 100ml bottles during early morning hours between 8.am to 10.am. The collected samples were preserved in 4% formalin.

The collected samples were placed in sedwig Rafter cell slide. The volume of cell 1/cm³ or 1ml exactly 1ml of sample is kept on the slide and a special cover glass is kept over it after organism settle. The chydoridae family order Alona were separated and again identified. The identification of the species were done with the help of standard key methods used for identification of cladocera with reference to chydoridae family.

Body parts like head, antennae, upper lip, labrum, rostrum beak like structure between antennules, post abdomen and coiled intestine are taxonomically important for identification of the chydoridae family species.



Result

Total 17 species have photographed and recorded from

both Nira left bank canal and Tarangawadi lake it seems that both the water bodies are rich source of cladocera (chydoridae) species it is good food source for pisciculture

shardabai Pawar Mahila Mahavidyalaya Barmati & Arts Science & Commerce college Indapur for providing infrastructural facilities and Laboratory equipment at Zoology Department

Acknowledgement

The author are thankful to Agriculture Development Trust

Sr	Species	Dorsal /Ventral	Antennules	Labrum	Postabdomen
1	Alona exigera	round	long	pointed (beak)	Trigonal with denticles
				And claw	
2	Pleuroxus				
	Trigonellus	Dorsal hood like	short	short	Elongated with pair of claw
3	Alonella				
	Rectangula	round (half moon shape)	elongated	pointed	elongated with claw
4	Chydorus				
	Sphaericus	round	short	beak like	elongated with denticles
				With single claw	
5	Ephemeroprus	Square shape	Short	pointed beak	present
6	Bosmina Longirostis	dor(round)	short	elephant trunk	
		Ven(vase shp)	like elongated	single claw	point
				spine at base	
7	Bosminopsis	dor(round)	short	elephant trunk	pair of claw
	Dieterus	vent(vase shp)		like elongated	
8	Alona Excusa	round	short	pointed curved	inside elongated dot on abdomen
9	Ephemeroporus	dor(round)	blunt	short	elongated with claw
	Orientalis	ven(square)			
10	C.Barroisi	Spherical	short	short stick like	pointed bears a
				basal spine	
11	Biapertura	round	Blunt	short	short with claw
	Affinis				
12	Acroperus Harpae	round	long	bifurcated	short with claw
			Beak like		
13	Alona Rustica	round	elongated	stick like	Triangular With
				Bifurcated	claw
			hairy		
14	Alonella	round	long	stick like	short with claw
	Globusa				
15	Alona	round	long	pointed	
	Combouei				
16	Dunhevedia Crassa	round	short	triangular shp	
17	Pleroxus similis	round	v short	short	Body transparent
				bifurcated	whitish in colour

Discussion

Chydoridae bears antennae situated in the head region is represented by formula 1-1-30-0-3 or 0-1-3 the antennae serves for locomotary organ. The upper lip labrum is taxonomically 0-0-1 important in chydoridae and it bears a keel. The presence or absence of the hair or setae and their size on the legs are counted for systematic position. The post abdomen may be short and stout it bear spine and denticles of various shape and arrangement are of great taxonomic important and the intestine is coiled in chydoridae. Rostrum is stout beak like structure situated between the antennules. It is long and pointed its shape and size are taxonomically important. In alona spp the ocellus is bigger than the eye brain optic ganglion and anterior part of the digestive tract in the head region. A pair of club shaped or slender antennules bearing sensory setae at the free end in the head region.

Both the water bodies of nira left bank canal and Tarangawadi are rich source of food to fishes which will be useful in increasing pisciculture good for production of carps.

References

- Brooks J.L.(1959) In fresh water biology, Eds ward , H.B. AND WHIPPLE , G.C John wiley & sons , inc London pp 587-656.
- Birge ,E.A (1918) The water fleas (cladocera) In fresh water biology editor , ward , H.B.and whipple, G.C.John wiley sons , inc new York
- Dadyay , E (1898) Microscopische suswasserthiera aus Ceylon termeszctr . fuz (Budapest) 21. 1-23.
- G.T.Tonapi Fresh water animals of India An Ecological approach
- Michael, R.G.(1973) Cladocera . in A guide to the fresh water organism J Madurai univ supp 1:1:2
- Nayar C.K.G.(1971) Cladocera of rajasthan , hydrobiology 37(3-4) 509 -519
- Patil , C S and Gouder B.Y. M.1982 fresh water fauna of dharwad (Karnataka state india) J Karnataka university 27 :115-126.
- Pennak .R.W.(1953) Fresh water invertebrates of the united states the Ronald press company, new york.
- Swammerdan, 1769 In fresh water biology .eds ward , H.B. and Whipple, G.C.John wiley and sons .inc new York
- Smirnov , N.N.1974 . Fauna of the ussr crustacean chydoridae vol Israel programme for scientific translation Jerusalem pp.1-644 (in English)