



Impacts of climate change on vector-borne diseases: a review of 2011 dengue outbreak in Pakistan

Saadullah Ayaz^{1*} and Abneer Anees²

¹IUCN, International Union for Conservation of Nature and Natural Resources, Pakistan.

²Environmental Sciences, COMSATS Institute for Information Technology, Islamabad.

ARTICLE INFO

Article history:

Received: 20 February 2013;

Received in revised form:

15 September 2014;

Accepted: 25 September 2014;

Keywords

Climate,
Vector-borne,
Diseases.

ABSTRACT

Dengue (dengue hemorrhagic fever) has emerged as an epidemic in Pakistan, since 1994. The recent dengue outbreak in year-2011 has drawn serious attention, which has affected over 16,000 people, out of which 306 are reported to have died. By analyzing the available scientific research, this paper reviews the consequences, which have resulted in increased activity of vector, i.e. 'Aedes aegypti' during year-2011. The paper establishes a link of the 2011 dengue outbreak with hydro-meteorological events, which have occurred over the year- 2011, which are attributed to climate change. The paper concludes that suitable climatic conditions prevailed during the year, which promoted the the insect's survival time and habitat but also its replication and maturation, leading to un-president epidemic outbreak of dengue. The paper also proposes appropriate policy planning and institutional, technical, operational/administrative and social and recommendations to manage and control any future dengue outbreaks in Pakistan.

© 2014 Elixir All rights reserved.

Introduction

Dengue is a acute febrile viral disease, that has drawn serious attention in recent decades. Dengue is caused by 'dengue fever virus (DENV)' which is a ribo-nucleic acid (RNA) virus of the family 'Flaviviridae' belonging to genus 'Flavivirus'. This virus is transmitted to humans by mosquitoes belonging to genus 'Aedes', which are widely distributed in sub-tropical and tropical regions of the World. Dengue is classified as a 'major global health threat' and nearly 2.5 billion people (two fifths of the World's population) is now at risk from dengue. The disease is endemic in more than 100 countries in Africa, the Americas, the Eastern Mediterranean, South-east Asia and the Western Pacific. The countries in South-east Asia and Western Pacific region are the most seriously affected because of the tropical climate and poor living conditions of these regions (WHO, 2009).

The symptoms of dengue virus include; fever, headache, muscle and joint pains (due to which it is also sometime referred to as 'break-bone fever'), severe eye pain (behind eyes), skin rashes (maculopapular rash) and severe itching. During dengue fever, hemorrhagic complications may also appear, such as bleeding from the gums, nosebleeds, and bruising (Lloyd, 2003). In some cases, the disease develops into the life-threatening 'dengue hemorrhagic fever (DHF)', resulting in bleeding, low levels of blood platelets and blood plasma leakage, or may lead into 'dengue shock syndrome', where dangerously low blood pressure associated by circulatory failure occurs. (CDC: Dengue).

Reviewing the trends and events, this paper provides an overview of the year-2011 dengue outbreak in Pakistan and attempts to establish a relationship of this incidence with the global phenomenon of climate change.

Epidemiology:

Dengue/ dengue hemorrhagic fever is present around the World, predominantly in urban and semi-urban areas. Several factors have combined to produce epidemiological conditions in developing countries (tropics and sub-tropics), that favor viral transmission by the vector, i.e. *Aedes aegypti*. Inadequate basic urban infrastructure (e.g. un-reliable water supply that leads householders to store water in containers close to homes) and increase in volume of solid waste, such as discarded plastic containers and other abandoned items, which provide dengue larval habitat for in urban areas, thus provides conducive conditions for dengue outbreak (Dengue: Guidelines for Diagnosis, Treatment, Prevention and Control, 2009)

Geographical expansion of the mosquito has been aided by international commercial trade particularly; in used tyres, which easily accumulate rainwater and remain un-noticed. Increased air travel and breakdown of vector control measures (bio-safety protocols) have also contributed greatly for dengue.

Transmission:

Dengue virus is transmitted to humans through the bites of infective female *Aedes* mosquito, particularly by '*A. aegypti*', which is one of the most efficient vectors, because it is highly anthropophilic and thrives in close proximity to humans. Other disease transmitting species include; '*A. albopictus*', '*A. polynesiensis*' and '*A. scutellaris*'. All of these species have a particular ecology, behavior and geographical distribution. Once infected, humans become the main carriers and multipliers of the virus, serving as a source of the virus for un-infected mosquitoes. The virus circulates in the blood of an infected person for 2-7 days after extrinsic incubation. At approximately the same time, the person starts developing symptoms of fever. Patients who are already infected with the dengue virus can transmit the infection via female *Aedes* mosquitoes, as soon as

the first symptoms appear. (WHO, Programmes and projects: Dengue)

Climate change and Dengue:

Climate Change is rapidly unfolding environmental and developmental challenges for the World, most of which have multi-natured catastrophic dimensions. Pakistan is amongst the most vulnerable nations and has been rated as 'at extreme risk' by a Climate Change Vulnerability Index (CVI), which has ranked Pakistan at number 16th in the list of 'most vulnerable countries of the World' over the next 30 years (Maplecroft, 2010).

Global warming threatens to further exacerbate the spread of many infectious diseases because increases in heat, precipitation and humidity, which can foster better conditions for tropical and subtropical insects to survive and thrive in places previously in-hospitable to those diseases. Warmer temperatures boost the speed of development of adult mosquitoes, increasing their numbers. Female mosquitoes bite more frequently in hotter temperatures and warmer winters enable mosquitoes to survive in areas that were formerly too cold. Higher temperatures also shorten the time it takes for the virus inside the mosquito to develop and become infective.

The effects of global warming on human health, besides others are visible in form of vector-borne infectious diseases such as dengue fever is mainly attributed to the expansion of the infested areas of vector (mosquitoes) that provides conducive conditions for increasing their number and also expansion of feeding activity. Changes in climate, that can affect the transmission of vector-borne infectious diseases include; temperature, humidity and altered rainfall (Ichiro, 2010).

Objectives:

The objectives of this review paper are;

- i. Conducting a desk review of the circumstances, which have resulted in dengue outbreak in Pakistan, during year-2011,
- ii. Through available information, establish a relationship of this incidence with the phenomenon of climate change,
- iii. Propose appropriate policy planning and institutional, scientific/technical, operational/administrative and social recommendations to control and/or better manage any future dengue outbreaks in Pakistan.

Review Of 2011- Dengue Outbreak In Pakistan

Over the past few years, Pakistan has emerged as a region of endemic dengue activity. The first outbreak of dengue fever (DF) was reported in Karachi in year-1994. Later, another outbreak emerged in the upper regions of Punjab in year-2003. There have been regular epidemics, with an increasing number of dengue cases in Karachi from year-2005 onwards.

The 2011 dengue outbreak in Pakistan, especially across Lahore and other cities of Punjab, including; Faisalabad, Sheikhpura, Rawalpindi and Multan etc., have drawn serious attention. A total of 16,000 people have been affected by October, 2011, out of which more than 306 have died. Highest number reported cases were from Lahore that is nearly 14000 (Government of Punjab, 2011).

Monsoon season, hot and humid climate favors mosquitoes breeding. This is main reason epidemics of dengue tend to coincide with the rainy season including; that in Pakistan (Shamim, 2010). Sub-sequent floods across the country, has indirectly lead to an increase in dengue, through the expansion in the number and range of habitats, like standing water caused by rainfall.

In dengue, an increase in temperature not only affects the insect's survival time and habitat, but also its replication, maturation and infective periods by providing suitable

conditions for mosquito survival (Lounibos *et al.* 2002). An overall warmer trend observed in different parts around the country has allowed these vectors to survive over winter, migrate into areas previously free of disease, or to trigger exacerbation of transmission in endemic areas.

Global warming is anticipated to increase the amplitude of the El Niño/southern oscillation (ENSO), resulting in marked changes in precipitation patterns. High frequency of rainfall events ensures that breeding places like; abandoned automobile tires, scupper drains, pots, buckets, cans and stagnant water in potted plants and the trays underneath them and also garbage dumps etc. creates larval habitats for mosquito, thereby expanding adult mosquito population (Cayan, 1999).

Studies show that the optimal temperature for dengue transmission is over 20°Celsius (°C). At temperatures below 20°C (68°Fahrenheit), the dengue mosquito cannot complete its growth cycle. Whereas temperature range below than 20°C up to freezing are unfavorable for mosquitoes. Contrary to prevailing suitable conditions during 2011, an extremely hot ambient temperature results in mortality of mosquitoes, while cold climate destroys larvae and eggs.

Pakistan Meteorological Department reported 10% above normal rainfall in 2011 in northern half of Pakistan including; Kashmir, Punjab and Khyber-Pakhtunkhwa provinces, at occasions by the interaction of easterly and westerly systems, resulting in heavy downpour in these areas (Pakistan Meteorological Department, 2010). Average for the monsoon rainfall in Lahore is 470.1 millimeters (18.51 inches) but in 2011; a total of 1,439.8 millimeters (56.69 inches) was recorded (Pakistan Meteorological Department, 2011), which is well above the yearly average.

In addition to climatic factors, the incidence of dengue outbreak in Pakistan is influenced by many demographic and societal factors like population growth, un-planned urban areas and inadequate public-health systems and in-appropriate facilities for handling and storing water.

Results And Discussion

The incidence of 2011 dengue outbreak across Pakistan is primarily attributed to global climate change, which has intensified the epidemic by providing favorable conditions for breeding of vector mosquitoes. Their transmission is also climate sensitive, as mosquitoes require standing water to breed. Humid conditions, higher rainfall the degree of urbanization were found to correlate with increasing risk of dengue fever, which generally prevailed in central Punjab during May-October, 2011 (Wu *et al.* 2009).

A favorable climate necessary for disease transmission is moist and moderately warm, has prevailed during 2011 in Pakistan. Other factors for an epidemic to trigger were, i.e. (i) an increase in vector populations, poor preparedness and susceptible human population, (ii.) Poor urban planning and other societal factors including; practices to store of water in open containers, (iii.) lack of public health infrastructure and awareness among people and capacity to adapt is weakest, which has contributed to emergence of dengue in year-2011.

Thus keeping in view all the above factors, it is concluded that in year-2011, conducive climatic conditions for the dengue vector prevailed over the Pakistan, which resulted in disease outbreak, particularly across Punjab province. If a comprehensive plan for prevention and control is not followed, dengue and other vector borne disease would continue to be a problem in days to come, in a scenario where climate change and other associated changes in weather patterns are anticipated to be more pronounced.

Recommendations

The following set of policy, technical, operational/administrative and social and recommendations are put-forth to control and/or manage any future dengue outbreak in Pakistan:

Policy Planning and Institutional Coordination Measures

- i. Sound institutional preparedness under clearly devised policy guidance is required, both at federal and provincial level. In order to control and/or manage the vector outbreak, clearly defined roles and needs to be devised and institutional responsibilities must be fixed. Sound institutional coordination mechanisms must also be put in place.
- ii. There is a need to further strengthen Pakistan's role in climate change diplomacy, through projecting incident like 2011-dengue outbreak. This would project Pakistan's stance as one of the 'most vulnerable nations', thus influencing the global dialogue.
- iii. Active participation of inter-sectoral partners is needed, right from the programme planning to implementation and evaluation, for strengthening the policy response. To deal with the health effects of the disease, the health sector must be engaged in preparedness activities at all levels; international, regional, national, local and community.
- iv. Short to medium-term (up to one year) and long-term (from one up to 5 years) multi-sectoral strategic plans must be devised at provincial level, with sufficient allocation of financial and administrative resources to adopt a comprehensive approach, covering; public health and safety, administrative and environmental measures.
- v. Integrating climate change concerns in health related policies and action plan at national as well as provincial level is essential. This would enable set priorities for mobilizing requisite capacities and resources.
- vi. Efficient and effective policy implementation and feedback/monitoring mechanisms must be put in place to ensure implementation of the policy measures and also ensure effective coordination amongst all relevant institutions/bodies and maintain federal to provincial coordination.

Scientific/Technical and Administrative Measures

- i. High-class collaborative scientific research on the epidemiology of vector and other factors, such as climatology needs to be conducted, to further understand exact dynamics of the vector and factors that influence such out-breaks. Policy studies backed by past, current and forecasted climate change conditions with historical epidemics must be conducted, in order to reveal the risk of dengue in days to come. The results of such research could supplement effective decision-making by supporting action planning.
- ii. Research on promotion of naturally occurring or introduction of new biological control agents for mosquitoes (parasitoids) needs to be adopted for controlling the vector. The carefully selected measures may include; importation, augmentation, and/or habitat management for parasitoids.
- iii. Epidemiological and entomological surveillance, outbreak investigations and meteorological forecast to anticipate climatological conditions over the year must be done to devise and support implementation of decision-making and ground-action. Field based research, complimented by use of tools such as Geographic Information Systems (GIS) must be used to keep a track of any incidence. Such data and information would help in devising early warning systems.

Ongoing local research is also needed, because global warming's effects on temperature and rainfall patterns can affect the risks of dengue outbreaks differently in specific areas.

- iv. Institutional strengthening all all tiers is necessary to effectively implement policy and programmes. Particularly, district administration in all 'high-vulnerable' districts must be capacitated to take necessary administrative/pre-emptive measures in this regard.
- v. Institutional strengthening of research facilities and and institutions related to outbreak response, including; hospitals must be strengthened. Necessary facilities including research equipment, medicines and others must be made available to manage any outbreak in more effective manner. Human capacities in diagnostic and analytical methods, needs to be build with comprehensive training programmes, including; those for scientists and medical staff (doctors and paramedics).
- vi. Standard case management is critical for reducing morbidity and preventing mortality due to dengue. Adoption of guidelines for clinically parasitology and needs to be followed, since transfusion of platelets in most of the cases is critical for saving lives, availability of safe blood products warrant strengthening the capacity of blood transfusion services to respond effectively during outbreaks.
- vii. Strict guidelines need to be set in place for water storage and disposal, particularly in urban and per-urban areas. Efficient enforcement mechanisms needs to be adopted for keeping a check on water storage facilities/tanks, ponds, fountains etc., which can offer breeding grounds for *A. aegypti*.
- viii. The use of chemicals for fogging and fumigation must be strictly regulated, as their negative impacts are often greater than what is intended. In-discriminate use may be a 'populous demand' and politically leveraged, but inappropriate use can counterproductively increase pest resistance. Also, studies suggest that very less of applied chemical is effective in eradication of mosquitoes, the rest reach destination killing non-target species (may be biological control agent) and contaminates air, water, food, individuals humans and soil.

Social Measures

- i. Human behavior and practice, which leads to breeding of mosquito vector needs to be changed, through continuous advocacy. Such targeted campaigns must be conducted which have high impact. Such campaigns should targeting women folks, school children, service providing institutions at local levels and public at large. Role of mass media is also crucial to rope in additional support. Involvement of local institutions (youth/welfare associations etc.) in campaigning, could also help in identification and implementation of locally acceptable measures related to water handling/storage, waste and sewage disposal.
- ii. Advocacy to control indoor mosquitoes population is also must for promotion of use of mosquitoes nets, and insect killing sprays and repellent like liquids/coils needs to be selectively promoted. Measures to minimize and/or limit outdoor activities particularly, at peek times of vector activity (dawn and dusk in case of dengue) should be practiced. Schools and other institutions needs to be particularly focused to take necessary measures in this regard.
- iii. Advocacy on the impacts of climate change on the country, particularly in health sector would be instrumental for long-term campaigning against the epidemics, such as dengue

References:

- Cayan D R, Redmond K T and Riddle L G (1999). ENSO and hydrologic extremes in the Western United States. *J. Climate* 12, 2881– 2893.
<http://journals.ametsoc.org/doi/pdf/10.1175/15200442%281999%29012%3C2881%3AEAHEIT%3E2.0.CO%3B2>
- Centers for Disease Control and Prevention (CDC): Dengue. <http://www.cdc.gov/Dengue/epidemiology/index.html>
- Dengue: Guidelines for Diagnosis, Treatment, Prevention and Control (2009): World Health Organization (WHO) and the Special Programme for Research and Training in Tropical Diseases (TDR) http://whqlibdoc.who.int/publications/2009/9789241547871_eng.pdf
- Government of Punjab (Pakistan). Combating Dengue. http://dengue.punjab.gov.pk/?q=system/files/Summary_12_Oct.pdf
- Ichiro Kurane MD (2010). The Effect of Global Warming on Infectious diseases. *Public Health Research Perspective*. Volume 1, Issue 1, Page 4-9
<http://www.kcdc-phrp.com/article/S2210-9099%2810%2900005-6/fulltext>
- Lounibos L P , Suárez S , Menéndez Z , Nishimura N , Escher R.L , Connell S M O and Rey J R (2002). Does temperature affect the outcome of larval competition between *Aedes aegypti* and *Aedes albopictus*? *Journal of Vector Ecology* 27(1):86–95. <http://dhf.ddc.moph.go.th/abstract/s11.pdf>
- Lloyd S L (2003). Best Practices for Dengue Prevention and Control in the Americas. Prepared for the USAID Bureau for Latin America and Caribbean under EHP Project 26568/E.V.4.LACDENGUE
http://www.ehproject.org/PDF/Strategic_papers/SR7-BestPractice.pdf
- Pakistan Meteorological Department (2010). National Weather Forecasting Center, <http://pakmet.com.pk/MON&TC/Monsoon/monsoon%282010%29.html>
- Pakistan Meteorological Department(2011) http://www.pakmet.com.pk/FFD/index_files/daily/rainfallsep1.htm
- Maplecroft (2010). Big Economies of the Future-Bangladesh, India, Philippines, Vietnam and Pakistan-most at risk from climate change
<http://maplecroft.com/about/news/ccvi.html>
- Shamim M (2010). Frequency, Pattern and Management of Acute Abdomen in Dengue Fever in Karachi, Pakistan. *Asian Journal of surgery*, Vol 33 No. 3
<http://linkinghub.elsevier.com/retrieve/pii/S101595841060019X?via=sd&cc=y>
- WHO Programmes and projects: Dengue. <http://www.who.int/denguecontrol/human/en/index.html>
- WHO (2009) Dengue and dengue hemorrhagic fever, Media center, Fact sheet
<http://www.who.int/mediacentre/factsheets/fs117/en/>
- Wu P C , Lay J G , Guo H R , Lin C Y, Lung S C, and Su H J (2009). Higher temperature and urbanization affect the spatial patterns of dengue fever transmission in subtropical Taiwan,” *Science of the Total Environment*, vol. 407, no. 7, pp. 2224–2233, 2009.