



Bio-Monitoring of metal deposition in ranthambhore national park (Rajasthan), India using *Riccia aravalliensis* pande et udar

Afroz Alam

Department of Bioscience and Biotechnology, Banasthali University, Banasthali, Rajasthan.

ARTICLE INFO

Article history:

Received: 8 October 2013;

Received in revised form:

19 March 2014;

Accepted: 29 March 2014;

Keywords

Bio-monitoring,
Elemental concentration,
Liverwort,
Riccia,
Ranthambhore.

ABSTRACT

In the present study the level of S, Cu, Cd, Zn, Pb and Cr was estimated in samples of the common thaloid liverwort *Riccia aravalliensis* Pande et Udar, from Ranthambhore National park, Rajasthan (India). High metallic load was observed both in substrate as well as in plant tissue at locations adjacent to higher vehicular load, during winter the metallic content is highest, followed by summer and monsoon season. Elemental concentration in substrate for *R. aravalliensis* was in the order of Zn > S > Pb > Cu > Cd > Cr, while in plant tissue it showed S > Zn > Pb > Cu > Cd > Cr which is indicative of air borne trace elemental load. Quantitative analysis of these elements in vegetative thalli of *R. aravalliensis* and the substrate showed an increase in metallic content during winter, which reveals the significance of this liverwort as a biological sink of mineral elements present in the soil and air and may provide an important tool in estimation of both aerial pollution and mineral enrichment in soil.

© 2014 Elixir All rights reserved

Introduction

In recent past, there has been a surging curiosity regarding bryophyte responses to environment due to their characteristic thallus composition as they are composed of sheets of tissue, therefore they are in close vicinity with the environment and respond more rapidly to environmental changes than vascular plants, because they get water, nutrients and elements mainly from the atmosphere. Richards [1] remarked that “the responses of bryophytes to edaphic factors are certainly sharper and clearer than the flowering plants”. Bryophytes are major constituent of hilly forest cover and pioneer in colonization. They play a vital role in carbon assimilation, mineral and moisture enrichment as well as contribute organic matter, nutrients materials and moisture to substratum which they absorb from air.

They are very handy tool in biological monitoring where the use of organisms provides information on certain aspect of biosphere like air-borne metal content. They are suitable and inexpensive method to quantify the possible emanation of metals because they are able to mount up elements in extremely high concentration and this provide easy detection of elements present even in very low concentration in the environment. On being trapped metals bind to the cell wall and remain retained for years and can be analyzed in future also. The lack of transportation system, results absorbed minerals to be adequately preserved.

Bryophytes tolerance and well suited and safe for the purpose of bio-monitoring [2] like other bio-monitoring organism they also possess the ability to survive quantity specific essential and non essential elements present in the environment and utilize them in their metabolism, these elements show toxicity at higher concentration by sensitive species, however, excess elements are detoxified by some molecular proteins in tolerant species. The work carried out earlier has been largely associated with musci as compared to hepaticae. Among the hepaticae, Marchantialean forms are considered to be most resistant against all types of adverse conditions as they are also called “resurrection plants” hence can

profitably be used for such studies. Several contributions mostly from abroad and elsewhere have been made in this direction [1-18]. But in India these type of studies is restricted to few workers only [19-22].

Material studied:

Considering the need of such studies it was thought desirable to assess the terrestrial liverwort (*Riccia aravalliensis*) of order Marchantiales as an indicator plant as well as accumulator of mineral elements/ heavy metals from soil and atmosphere.

Characteristics of *Riccia aravalliensis*:

Thallus green, large, dichotomously branched, apex notched. Midrib somewhat indistinct. Ventral surface covered with numerous rhizoids and reduced hyaline scales in single row on each side of midrib. Thallus internally well differentiated into upper assimilatory zone having air chambers in single row with linear assimilatory filaments, storage zone present up to the margins.

Studied Area:

The Ranthambhore Tiger Reserve is the single biggest stretch of dry deciduous *Anogeissus pendula* forest left unbroken in India. In Ranthambore the biodiversity is made even richer by the invasion of the Vindhyan hill system. The terrain of Ranthambhore tiger reserve is mostly craggy and hilly and is intimately related to the Great boundary fault. The highest point of this is Gazella peak, 507 m above sea level. The lowest altitude of this tract is 244 m above sea level at Bodal. Streams flowing in northern tract form the catchment of the river Banas and streams flowing in southern tract drain directly in the river Chambal. Its cardinal points are 25° 54' to 26° 12' N Lat. and 76° 23' to 76° 39' E Long., annual average rainfall 800m, temperature range 4°C (minimum) -47°C (maximum). The annual mean humidity is 72%, which reaches to its maximum (99%) during monsoon. The sand stone beds of these hills are flat-topped and form extensive table lands known as "Dangs". These dangs rise abruptly from flat ground and have sandstone

Tele:

E-mail addresses: afrozalamsafvi@gmail.com

© 2014 Elixir All rights reserved

ridges running continuously along their edges. At places, small and short-lived streams have eroded deep, long and narrow gorges that are locally known as "Khoahs" [23].

Material and Methods

The materials used for the present study include (i). The vegetative thalli of *Riccia aravalliensis* and, (ii). The soil samples of substrata from different sites on which they grow. Vegetative thalli of *Riccia aravalliensis* and soil samples from the substrate from 2 major regions (peripheral zone and undisturbed zone) of Ranthambhore National Park were collected between 15.05.2012 to 05.02.2013. The quantitative analysis of S, Fe and heavy metal content (S, Cu, Cd, Zn, Pb and Cr) was carried out in both the plant thalli and soil samples from different localities. The soil samples were also analyzed for its physical and chemical properties to find out the similarities and differences in soil properties of different regions. The presence of mineral elements such as S, Cu, Cd, Zn, Pb and Cr in soil as well as in the plant tissue was estimated to see the extent of accumulation of soil elements in plant material.

The plant material was collected along with soil samples from different sites and were individually sealed in polythene bags and brought to laboratory. The plants were examined under microscope and identified for their taxonomic status. The substrate matter attached to the plant material was removed by high pressure washing with tap water and then by repeated washing with distilled water and finally immersed in double distilled water for 2 hours. The meticulously washed plant samples employed for chemical analysis were dried at 70° C for 48 hours in labeled bamboo paper bags. The oven dried plant material was used for analysis of some macro and micronutrients (i.e., S, and Fe content) after wet digestion employing Piper's method [24]. Further estimations were done according to the methods employed by various workers as tissue Sulphur [25], and tissue Copper, Chromium, Cobalt and Iron were analyzed by DTPA extraction method [26] using Atomic Absorption Spectrophotometer (AAS).

The pH of soil was measured using digital pH meter, and the concerned elemental content estimated by specific methods employed by different workers for particular elements, such as available S [27].

Observations

Result and discussion:

Sulphur (S):

High influx of tourist, the consumption of fuels can be the reason for high increase in S content throughout the year near densely populated areas [28]. The result is clarifying the statement that at tourist places there will always be escalating drift observed for vehicle pollution; consequently increase in the metals, associated with the vehicular pollution [29, 30] as it appear as many fold increase in its value during winter could be due to enhanced traffic in winter seasons. The overall maximum percent (%) S load in location of high traffic movement at their maximum value (Table 1, 2; Fig. 1a-1c; Fig.2a-2c).




Moreover, the tourist influx during winter in this tourist spot find itself choked with tourist traffic in and indeed they are more crowded in each season except monsoon [31]. The deposition values are positively related to the traffic (and population) density. This reflects their high value in plants growing at petrol pumps or along the road (near taxi stands) in Ranthambhore compared to other catchment areas.

Comparison of metal S concentration in different seasons exhibited its lowest concentration in rainy season could be attributed to decrease of tourists in the rainy season and secondly pollutant leaching and increase in growth and biomass

occurs more rapidly as a result it reduces the sulphur percentage in leaf in proportion to biomass [32].

Lead (Pb):




Excessive fuel consumption could be the reason for its high value in all seasons near proximity to the town, along the roads or near these hill stations' taxi stand which intensified during winter, a tourist season (Halleraker et al., 1998). The result is explanatory the statement that at tourist places there will always be increasing trend observed for vehicle pollution; consequently increase in the metals, associated with the vehicular pollution [29]. Present experimental data further revealed that many fold increase in its value during winter could be due to enhanced traffic in winter seasons (Table 1). The overall maximum percent (%) metal Pb load in both locations stands with their maximum value (Table 1, 2; Fig. 1a-1c; Fig.21a-2c). A decrease in metal lead load during monsoon could be due to the atmospheric metal load is washed during heavy rains ((Table 1, 2; Fig. 1a-1c; Fig.2a-2c). Combustion of leaded, low-leaded and unleaded gasoline continues to be the major source of atmospheric Pb emissions [33]. A decrease in metal lead load during monsoon could be due to the atmospheric metal load is washed during heavy rains.



Since high-octane petrol (with 0.45 g lead/l) is still used in the majority of the cars, the deposition values are positively related to the traffic (and population) density. This reflects their

high value in plants growing at petrol pumps or along the road (near taxi stands) in as compared to other catchments areas.

Like Sulphur comparison of metal Pb concentration in different seasons revealed its lowest concentration in monsoon season due to decrease of tourists in the rainy season and leaching increase in growth and biomass occurs more rapidly as a result it reduces the metal percentage in thalli in proportion to biomass [32].



Zinc (Zn):

Maximum percent increase of metal Zn was observed at the same locations similar to Pb. A seasonal trend for metal Zn justifies that winter had high value of Zn followed by summer and rain (Table 1, 2; Fig. 1a-1c; Fig.2a-2c). Heavy rain fall could be one another way to explain the decrease in Zn load as most of the metal Zn was surface adsorbed [34]. The enrichment ratios for Zn in thalli were not constant throughout the year; therefore, interpretation of enrichment of Zn is complicated [35]. Here, the role of traffic cannot be over ruled to increase Zn content of the air as it is the part of automobile stratum [36].

Copper (Cu):

In comparison to Pb and Zn, a moderate deposition rate of metal copper was observed from same catchment areas. Metal Cu was high in populated as well as rural areas. Maximum value of copper was in peripheral zone during all the three seasons (Table 1, 2; Fig. 1a-1c; Fig.2a-2c). Therefore, present finding is further supported by the presence of abundant Cu absorbing bryophytes in these areas. A decrease in Cu in summer with respect to winter could be explained by considering that dry deposition increases on moving from humid to arid climates [37]. Concentration and distribution pattern of both metal Zn

and Cu in *Riccia aravalliensis* were quite similar and both metals were high in rural transplants located in vicinity of national park. Cu pollution may also originate from domestic waste disposal. The use of CuSO₄ mixed kerosene oil could also one of the facts of increase of Cu concentration in domestic areas.

Cadmium (Cd):

Cadmium metal is very easily leached out from surface and the similar results were observed in monsoon season in *Riccia aravalliensis*. All catchment's area shows significant decrease in metal Cd value in rainy season with respect to summer and winter (Table 1, 2; Fig. 1a-1c; Fig.2a-2c). Easily leaching property could be the reason for their low value even in the forest areas [33]. There is a frequent use of breaks in the hilly region compare to the land areas. This could be the reason for high value of Cd in peripheral zone during winter. An increase in the Cd on such places could be from abrasion of clutch, breaks of the vehicles. Higher concentration in agricultural land might be due to the use of phosphate fertilizers [38]. Perhaps Cd is also present in the petrol as mining impurities. Service shops related with metals are the other factors for this increase.

Percent metal load and pollution index value:

The pollution index value at peripheral zone and undisturbed zone of Ranthambhore National Park was +1.212453 and +0.877541 respectively.

The PI value was highly positive at the areas near by taxi stand and at petrol pumps. This was further supported by percent metal loading in respective locations during these years. The maximum positive values were measured in proximity to the city area. Negligible value at undisturbed zone reveals that it is relatively cleaner site (pollution free). The effect of intensity of traffic compared to the influence of other factors like farming on sampling points are much higher as the spots were nearer to the roads (except control site). Besides this, contamination was high and reduces with distance [39].

Discussion

The bryophytes in general are not only efficient absorbers of heavy metals but also useful indicators of aerial fallout of heavy metals usually much more than the vascular plants [40-41]. These studies have proven that the chemical analysis of carpet forming bryophytes is the rapid and comparatively economical tool for surveying heavy metal deposition in terrestrial ecosystem and in surrounding ambient air. The monitoring by bryophytes can be done even in any remote area where no expensive technical equipment can be transported for estimation. Collection and sampling is an easy task in case of bryophyte and can be done by any one suspecting that air has been contaminated. Their habitat diversity, structural simplicity, totipotency and rapid multiplication rate, make them an ideal organism for studies related to pollution and mineral enrichment besides phyto-sociological and eco-physiological aspects. The mineral elements (S, Pb, Zn, Co, Cr and Cu) estimated in thallus tissue of *Riccia aravalliensis* are not only essential elements but also contribute as an important component of the plant organization by playing major role in their metabolic activities. Heavy metals such as Cu, Co, and Cr have also been estimated to find out their extent of accumulation in the thalli from soil as well as aerial fall out. The sites of undisturbed zone have been considered as control as the conditions in this region is almost ideal for the growth of *Riccia aravalliensis* because of least population and least atmospheric pollution as compared to the sites in peripheral zone. Likewise, winter season is more affected with metals' pollution due to heavy influx of tourists to the national park from country and abroad.

Table 1: Metals concentration ($\mu\text{g/g}$ dry weight.) \pm SE in seasonally exposed samples of <i>Riccia aravalliensis</i> at two different localities of Ranthambhore National park. Significance differ from control (a) $p \leq 0.05$ and (b) $p \leq 0.01$ significance level						
Study sites	Winter (Fig. 1a)					
	Cu	Cr	Zn	Pb	S	Co
Peripheral (Disturbed Zone)	18.15 \pm 0.98	5.17 \pm 0.74	27.90 \pm 0.94	21.56 \pm 1.02	20.90 \pm 0.72	6.10 \pm 0.72
Central (Undisturbed Zone)	12.10 \pm 0.87	1.47 \pm 0.16	17.36 \pm 0.99	10.36 \pm 0.81	08.20 \pm 0.82	02.20 \pm 0.77
Summer (Fig. 1b)						
Peripheral (Disturbed Zone)	26.25 \pm 0.78	4.82 \pm 0.52	29.71 \pm 0.42	25.86 \pm 0.92	23.70 \pm 0.32	06.70 \pm 0.12
Central (Undisturbed Zone)	17.32 \pm 0.62	2.27 \pm 0.14	19.26 \pm 0.32	11.46 \pm 0.43	10.60 \pm 0.72	03.05 \pm 0.64
Monsoon (Fig. 1c)						
Peripheral (Disturbed Zone)	11.58 \pm 0.72	2.35 \pm 0.24	22.57 \pm 0.17	15.34 \pm 0.81	19.70 \pm 0.32	02.80 \pm 0.23
Central (Undisturbed Zone)	5.26 \pm 0.72	0.77 \pm 0.15	8.08 \pm 0.72	7.23 \pm 0.72	5.10 \pm 0.32	0 1.27 \pm 0.72

Table 2: Metals concentration ($\mu\text{g/g}$ dry weight.) \pm SE in soil samples at <i>Riccia aravalliensis</i> at two different localities of Ranthambhore National park. Significance differ from control (a) $p \leq 0.05$ and (b) $p \leq 0.01$ significance level						
Study sites	Winter (Fig. 2a)					
	Cu	Cr	Zn	Pb	S	Co
Peripheral (Disturbed Zone)	240.12 \pm 0.12	41.22 \pm 0.15	60.11 \pm 0.14	79.17 \pm 0.12	14.82 \pm 0.32	11.10 \pm 0.24
Central (Undisturbed Zone)	69.18 \pm 0.24	25.3 7 \pm 0.22	33.26 \pm 0.12	27.15 \pm 0.82	04.24 \pm 0.12	04.12 \pm 0.15
Summer (Fig. 2b)						
Peripheral (Disturbed Zone)	178.16 \pm 0.32	34.14 \pm 0.62	52.11 \pm 0.15	74.34 \pm 0.24	13.89 \pm 0.12	11.87 \pm 0.32
Central (Undisturbed Zone)	59.18 \pm 0.24	19.77 \pm 0.22	24.36 \pm 0.12	23.15 \pm 0.82	04.40 \pm 0.12	03.92 \pm 0.15
Monsoon (Fig. 2c)						
Peripheral (Disturbed Zone)	158.11 \pm 0.12	31.34 \pm 0.12	34.26 \pm 0.12	62.34 \pm 0.12	12.19 \pm 0.32	10.77 \pm 0.12
Central (Undisturbed Zone)	32.12 \pm 0.82	15.27 \pm 0.28	18.46 \pm 0.42	18.72 \pm 0.24	03.42 \pm 0.32	02.98 \pm 0.24

Conclusion

On the basis of present study it has been revealed that *Riccia aravalliensis* may be considered as accumulator of certain elements, whose concentration in plant tissue is generally related to the element content in the supporting substrate (soil). However, this relationship tends to vary in case of certain elements such as S due to aerial fall out in polluted areas. The tissue chemical content of the same species also varies according to its changed climatic conditions, but the basic level of requirement of a particular element is almost the same. The variation caused in the tissue chemical content may be due to the effect of pollutants (Table 1, 2; Fig. 1a-1c; Fig.2a-2c).

The Bryo-vegetation is an exploratory "tool" and should be widely accepted as readily as have rocks, soils, water and stream sediments. It should not be taken as a "nuisance" and must not be mercilessly bulldozed out because they accumulate pollutants and make the surrounding atmosphere free from the pollution (Brooks, 1972) and is evidently supported in the present investigation. The *Riccia aravalliensis* thalli accumulate S to a large extent besides serving as sink for heavy metals like Co, Cu, Pb, Zn and Cr and may act as soil purifier. Further investigations employing other species of bryophytes may profitably be used in monitoring environmental pollution as well as indicators of mineral enrichment of soil.

Acknowledgements

The author is grateful to Prof. Aditya Shastri, Vice Chancellor, Banasthali University, Rajasthan (India) and Prof. Vinay Sharma, Dean, faculty of Science and Technology, Banasthali University, Rajasthan for providing necessary support.

Reference

- Richards PW. Ecology In: Verdoorn, F., (Ed.). Manual of Bryology, 1932, 367-395.
- Giordano S, Admo P, Sorbo S and Vingiani S. Atmospheric trace metal pollution in the Naples Urban Area based on results

from Moss and Lichen Bags. Environ. Pollut, 2005, 136: 431-442.

- Pearson H. Studies in "copper mosses". J. Hattori Bot. Lab, 1956, 17: 1-18.
- Martensson, D and Berggren A. Some notes on the ecology of "copper mosses". Oikos, 1951, 5: 99-100.
- Schatz A. Speculation on the ecology and photosynthetic of the copper mosses. Bryologist, 1955, 58: 113-120.
- Naguchi A. On some species of *Merceya*, with special reference to variation and ecology; Kumamoto, J. Sci, 1956, Ser. B, Sect. 2(2): 239-257.
- Naguchi A and Furato H. Germination of spores and regeneration of leaves of *Merceya ligulata* and *M. gedeana*. J. Hattori Bot. Lab, 1956, 17: 32-44.
- Canon HL. Botanical prospecting for ore deposits. Science, 1960, 132: 592-598.
- Shacklette HT. The use of aquatic bryophytes in prospecting. Jour. Geochem. Expl, 1984, 21: 89-93.
- Brooks RR. Bryophytes as a guide to mineralization. New Zeal. J. Bot, 1972, 9: 674-678.
- James PW. The effect of air pollutants other than hydrogen fluoride and sulphur di oxide on lichens. In: Ferry B.W., Baddeley M.S., Hawksworth D.L. (Eds.), Air pollution and Lichen. University of Toronto Press, 1973, 143-175.
- Crundwell AC. *Ditrichum plumbicola* a new species from lead mine waste. J. Bryol, 1976, 9: 167-169.
- Rejment GI. Concentration of heavy metals, lead, Iron, manganese, zinc and copper in mosses. J. Hattori Bot. Lab, 1976, 41: 225-230.
- Ward NI, Brooks RR and Reeves RD. Copper, cadmium, lead and zinc in soils, streams, sediments, water and natural vegetation around the Tuimine, Te Aroha, New Zealand. N.Z. Jour. Sci, 1976, 19: 1981-1989.
- Shacklette HT and Erdman JA. Uranium in spring water and bryophytes at Basin Greek in Central Idaho. Jour. Geochem. Expl, 1982, 17: 89-93.

16. Smith SC. Base metal and mercury in bryophytes and stream sediments from a geochemical reconnaissance survey of Chadulur Quadrangle, Alaska. *J. Geoch. Expl.*, 1986, 25: 345-365.
17. Satake KA. "Copper moss" *Scopelophila cataractai* and copper accumulation. *Proc. Bryol. Soc. Japan*, 1951, 5: 105-107.
18. Shaw J and Schneider RE. Genetic biogeography of the rare "copper moss" *Mielichhoferia elongata* (Bryaceae). *Amr. J. Bot.*, 1995, 82: 8-17.
19. Saxena DK. Schatz, A., 1955. Speculation on the ecology and photosynthetic of the copper mosses. *Bryologist*, 1982, 58: 113-120.
20. Pant G. and Tewari SD. Bryophytes as Biogeoindicators: Bryophytic Associations of Mineralenriched substrates in Kumaon Himalaya In: *Topics in Bryology*, 1998, R.N. Chopra (ed.) Allied Publish. Ltd.
21. Chopra RN and Kumra PK. *Biology of bryophytes*, 1991, Wiley Eastern Ltd. New Delhi.
22. Glime JM and Saxena DK. *Uses of Bryophytes*, 1991, Today and Tomorrow Printers and Publishers India.
23. Alam A, Sharma V and Sharma SC *Bryoflora of Ranthambhore Tiger Reserve, Rajasthan (India)*, *Arch. Bryol.*, 2011, 106: 1-8. (2011).
24. Piper CS. *Soils and plant analysis*, Waite Agric. Res. Inst. The Univ. of Adelaide Australia, 1942, 251- 257.
25. Chesnin L and Yein CH. Turbid metric determination of available sulphate. *Proc. Soil Sci. Amer*, 1951, 15: 149-151.
26. Lindsay WL and Norvell WA. Development of DTPA soil testing for Zinc, Iron, Maganese and Copper. *Soil Sci. Soc. Am. J.*, 1978, 42: 421-428.
27. Olsen SR, Cole CV, Watanbe FS and Dean LA. Estimation of available phosphorus in soil by extraction with sodium bi carbonate, 1954, U.S.D.A. 939.
28. Thöni L, Schnyder N and Krieg F. Comparison of metal concentrations in three species of mosses and metal freights in bulk precipitations. *Fresenius' J. Analytical Chem.*, 1996, 354: 704-708.
29. Poikolainen J, Kubin E, Piispanen J and Karhu J. Atmospheric heavy metal deposition in Finland during 1985-2000 using mosses as bioindicators. *Sci. total. Environ.*, 2004, 318: 171-185.
30. Alam A, Srivastava SC. *Marchantia paleacea* Bert. As an indicator of heavy metal pollution. *Indian Journal of Forestry*, 2009, 32(3): 465-470.
31. Saxena DK and Saxena A. Uptake of metals in *Plagiochasma* and their uses in pollution monitoring. *Geophytology*, 2000, 28(1& 2): 129-137.
32. Saxena DK, Tuba Z and Arfeen MS. Seasonal passive metal monitoring during year 2003 to 2006 in Nainital of Kumaon hills (INDIA) by moss *Racomitrium crispulum*. *Acta Botanica Hungarica*, 2010, 52(1-2): 273-297.
33. Fernández JA, Aboal JR and Carballeira A. Testing differences in methods of preparing moss samples. Effect of washing on *Pseudoscleropodium purum*. *Environ. Monit. Assess.*, 2009, ISSN 1573-2959 (Online).
34. Bate JW. Mineral nutrient acquisition and retention by bryophytes. *J. Bryol.*, 1992, 17:223–240.
35. Makhholm MM and Miladenoff David J. Efficacy of a biomonitoring (moss bag) technique for determining element deposition trends on a mid – range (37 Km) scale. *Environmental Monitoring and Assessment*, 2005, 104: 1-18.
36. Shahmansouri MR, Pourmoghadas H, Parvaresh AR and Alidadi H. Heavy Metals Bioaccumulation by Iranian and Australian Earthworms (*Eisenia fetida*) in the Sewage Sludge Vermi-composting. *Iran. J. Env. Health Sci. Eng.*, 2005, 2 (1): 28-32.
37. Couto JA, Aboal JR, Fernandez JA and Carballeira A. A new method for testing the sensitivity of active biomonitoring: an example of its application to a terrestrial moss. *Chemosphere*, 2004, 57(4): 303-308.
38. Otvos E, Pazmandi T and Tuba Z. First national survey of atmospheric heavy metal deposition in Hungary by the analysis of mosses. *Sci. Total. Environ.*, 2003, 309: 151-160.
39. Alam A and Sharma V. Seasonal variation in accumulation of heavy metals in *Lunularia cruciata* (Linn.) Dum. at Nilgiri hills, Western Ghats, *International Journal of Biological Science and Engineering*, 2012, 3 (2): 91-99.
40. Richards LA. Filter funnel's from soil extracts. *Agron. Jour*, 1949, 41: 466-469.
41. Brüning F and Kreeb KH. Mosses as Biomonitors of heavy metal contamination within urban areas. In: Markert, B. (Ed.) *Plants as Biomonitors: Indicators for Heavy metals in Terrestrial Environment*, 1993, VCH, Weinheim and New York.